arXiv:0705.3653v1 [cond-mat.str-el] 24 May 2007

Mott Fermionic "Quantum" Criticality Beyond Ginzburg-Landau-Wilson

Zaira Nazario* and David I. Santiago†
$\star$ *Max Planck Institute for the Physics of Complex systems, Nöthnitzer Str. 38, 01187 Dresden, Germany*
$\dagger$ *Instituut-Lorentz for Theoretical Physics, Universiteit Leiden,*
P. O. Box 9506, NL-2300 RA Leiden, The Netherlands

(Dated: February 1, 2008)

The Mott critical point between a metal and a correlated insulator has usually been studied via density or spin density bosonic mode fluctuations according to the standard Ginzburg-Landau-Wilson phase transition paradigm. A moment's reflection leads to increasing doubts that such an approach should work as the transition is nonmagnetic, voiding the relevance of spin density modes. Charge density modes are irrelevant since the long range Coulomb interaction leads to a large plasmon gap and their incompressibility. In solidarity with these doubts, recent measurements of the Mott critical point in low dimensional organic materials yield critical exponents in violent disagreement with the bosonic mode criticality lore. We propose that fermionic fluctuations control the behavior of the Mott transition. The transition thus has an intrinsic quantum aspect despite being a finite temperature phase transition. We develop this hitherto unexplored physics, obtain experimental predictions and find agreement with one of the novel unexplained experimental exponents. We conclude that this Mott transition corresponds to a new universality class of finite temperature critical points that contains quantum effects and cannot be accounted for by conventional Ginzburg-Landau-Wilson wisdom.

PACS numbers: 71.10.Fd, 71.10.Hf, 71.30.+h

Strong Mott correlation physics is at the center of some the most spectacular and important phenomena in condensed matter physics: quantum antiferromagnetism in low dimensions[1], high temperature superconductivity (high T_c cuprates)[2], heavy fermion behavior[3, 4], interesting phase competition and their concomitant quantum and classical phase transitions, etc. In particular, a very remarkable experiment was published recently[5]. The phase transition between a Mott insulating phase and a metallic phase was measured in the quasi-2D organic material κ-(BEDT-TTF)$_2$Cu[N(CN)$_2$]Cl (phase diagram in Figure 1). The importance of the measured properties of the Mott transition lie in its novel unexpected exotic nature.

The critical exponents measured in the experiment *are in violent disagreement with those obtained from any of the universality classes that follow from the Landau-Ginzburg-Wilson phase (LGW) transition paradigm[6, 7].* This conventional and usually successful LGW understanding of phase transitions gets critical properties from the fluctuations of order parameter bosonic modes. Even though the Mott transition occurs between a featureless correlated insulator and a featureless metal, interesting approximate mappings to the liquid-gas transition (Ising universality class) seem to lead to an LGW picture of the transition[8, 9] driven by particle density modes.

In the Mott system, there are reasons to think that if an LGW picture is to work, the particle density modes would be the main players. The system consists of the fermionic quasiparticles and their quasi-bound state spin and particle density modes. Since there is no magnetism in either of the phases, spin density modes play no rule. This seem to single out density modes as the only possible candidates for LGW physics for the Mott transition. On the other hand, complete disagreement of predicted exponents from measured ones[5] points to the irrelevance of density modes and toward the Mott transition being a completely novel universality class.



FIG. 1: Pressure-temperature phase diagram of the family of compounds κ-(BEDT-TTF)$_2$X studied by F. Kagawa, *et. al.* on reference [5]. The measurements focused on the member κ-(BEDT-TTF)$_2$Cu[N(CN)$_2$]Cl.

With the hindsight of the experimental disagreement with density mode LGW physics, we can find important physical reasons as to why the density modes should be irrelevant at the transition. In electronic systems, proper determination of the effects of the long range Coulomb interactions of mobile electrons in a background of very heavy positive ions leads to a large plasmon gap, and thus to incompressibility for long wavelength density modes, with a leftover short range interaction between the electron-like and hole-like quasiparticles in a metal[10]. Such a short range interaction justifies the neglect of the long range Coulomb interaction among quasiparticles as it has been canceled via screening effects. Neglect of such long range effects in the interactions between density modes is incorrect as the plasmon gap is large and prevents the long wavelength density modes from becoming soft and playing an important role in the transition.

On the face of the experimental irrelevance of spin and particle density bosonic modes as drivers of the transition, we identify the quasiparticle fermionic degrees of freedom and their fluctuations as the main actors driving and determining the Mott critical transition. The fermionic quasiparticles are the only remaining modes in the system[1]. that can play a role at the transition. We see that even though the Mott transition we are considering is a finite temperature transition, fermionic quasiparticles and thus their statistics, play an essential role at the transition.

More importantly fermionic fields cannot acquire a nonzero macroscopic average as no two of them can occupy the same quantum state; fermions cannot be order parameters. Thus despite being a finite temperature transition, the Mott transition contains an essential quantum aspect to it stemming from the relevance of fermionic quasiparticles and contains no order parameter physics. *Transitions of this type, finite temperature but with an essential quantum aspect to them, are completely novel, are not compatible with the Ginzburg-Landau-Wilson paradigm[6, 7].* They have hitherto never been studied.

In order to study more quantitatively the Mott transition, we start from the partition function for the Hubbard model:

$$
\begin{aligned}
Z &= \int \mathcal{D}\psi^\dagger_\sigma \mathcal{D}\psi_\sigma e^{-S} \\
S &= \int_0^\beta d\tau d^2x \left[\sum_\sigma \left(\psi^\dagger_\sigma \frac{\partial \psi_\sigma}{\partial \tau} - \frac{\mu}{\hbar} n_\sigma \right) \right. \\
&\quad \left. + \frac{\hbar}{2m} \sum_\sigma \nabla \psi^\dagger_\sigma \cdot \nabla \psi_\sigma + \frac{Ua^2}{\hbar} n_\uparrow n_\downarrow \right] \\
&= \int_0^\beta d\tau d^2x \left[\sum_\sigma \left(\psi^\dagger_\sigma \frac{\partial \psi_\sigma}{\partial \tau} - \frac{\mu}{\hbar} \psi^\dagger_\sigma \psi_\sigma \right) \right. \\
&\quad \left. + \frac{\hbar}{2m} \sum_\sigma \nabla \psi^\dagger_\sigma \cdot \nabla \psi_\sigma + \frac{Ua^2}{\hbar} \psi^\dagger_\uparrow \psi_\uparrow \psi^\dagger_\downarrow \psi_\downarrow \right] .
\end{aligned}
\tag{1}
$$

where $\sigma = \uparrow, \downarrow$ represents the spin and a is the lattice spacing. The Mott critical point will occur at critical values $U = U_c$, $T = T_c$. The Hubbard interaction is somewhat inconvenient to manage due to its quartic nature. It can be written more conveniently with the help of a Hubbard-Stratonovich decoupling leading to

$$
\begin{aligned}
Z &= \int \mathcal{D}\lambda \mathcal{D}\rho \mathcal{D}\sigma \mathcal{D}\psi^\dagger_\sigma \mathcal{D}\psi_\sigma e^{-S} \\
S &= \int_0^\beta d\tau d^2x \left[\sum_\sigma \psi^\dagger_\sigma \frac{\partial \psi_\sigma}{\partial \tau} - \frac{\mu}{\hbar} n \right. \\
&\quad + \frac{\hbar}{2m} \sum_\sigma \nabla \psi^\dagger_\sigma \cdot \nabla \psi_\sigma + \frac{Ua^2}{4\hbar} \sigma^2 + \frac{Ua^2}{4\hbar} \rho^2 \\
&\quad \left. + \frac{Ua^2}{2\hbar} s\sigma + i \frac{Ua^2}{2\hbar} n\rho \right] ,
\end{aligned}
\tag{2}
$$

where we have defined $n(\vec{x}) \equiv n_\uparrow(\vec{x}) + n_\downarrow(\vec{x})$ and $s(\vec{x}) \equiv n_\uparrow(\vec{x}) - n_\downarrow(\vec{x})$. The partition function is that of fermions interacting with bosonic fields that correspond in a sense to particle and spin density modes ($i\rho$ and σ). This is an exact partition function for the Hubbard model and gives the correct description of its different phases and phase transitions.

The Hubbard interaction generates an effective interaction between particle and holes and spin density and density modes, as shown diagrammatically in Figure 2, given by



FIG. 2: Feynman diagrams representing the interaction of fermions with spin density and density fluctuations. Straight lines represent fermions, zig-zag lines represent spin density fluctuations and wavy lines represent density fluctuations.

$$
\begin{aligned}
&- \frac{T}{\hbar} \sum_n \int \frac{d^2\vec{k}}{(2\pi)^2} \left[i g_\rho n(\omega_n, \vec{k}) \rho(-\omega_n, -\vec{k}) \right. \\
&\left. + \; g_\sigma s(\omega_n, \vec{k}) \sigma(-\omega_n, -\vec{k}) \right]
\end{aligned}
\tag{3}
$$

where $g_\rho = g_\sigma = (Ua^2)/(2\hbar) = g_{rs}$ and bosonic Matsubara frequency $\omega_n = 2\pi n T/\hbar$. The dynamics of the collective modes and their effective interactions with the quasiparticles are equivalent to, and take the place of, the Hubbard interaction. The Hubbard interaction between fermions generates also dynamics for density modes and spin density modes via fermionic fluctuations. In fact, particle-hole fluctuations of the form illustrated by the Feynman diagrams in Figure 3 generate a term in the Boltzmann weight of the partition function or effective "Euclidean action" S describing the dynamics of the short range density and spin density fluctuations induced by the onsite repulsion.

The effective theory of the system consists of metallic fermions, density and spin density modes, and the interaction between these collective modes and the fermions

---

[1] The density and spin density modes, while not driving the transition, do play an indirect but not unimportant role as they can mediate interactions between the fermionic quasiparticles. Without such an interaction, no transition would be possible.



FIG. 3: Fermion fluctuation responsible for generating density and spin density fluctuations dynamics. The solid line represents the fermions, the wavy lines are density fluctuations and the zig-zag lines are spin density fluctuations.

with strength g_{rs}. Single particle or hole fluctuations will dress or renormalize the interactions between the density modes and the fermions, either enhancing or diminishing them. Processes of the form shown in Figure 4 change the value of the effective coupling constant g_{rs}.



FIG. 4: Feynman diagrams representing the renormalization of the interaction of fermions with spin density and density fluctuations. Straight lines represent fermions, zig-zag lines are spin density fluctuations and wavy lines are density fluctuations.

When we consider large momenta or small distance fluctuations, i.e. those between the microscopic momentum cutoff Λ (taken to be the inverse lattice spacing $1/a$) and a lower cutoff μ, processes of this type give a renormalized coupling constant appropriate to the distance scale $1/\mu$,

$$g_{rs}^{\mu} = \frac{g_{rs}^{\Lambda}}{1 - \frac{4}{3}\pi^4 \, \left(g_{rs}^{\Lambda}\right)^2 \, \frac{\hbar^2 m v_F^2 \Lambda^2}{T^3} \, \Lambda(\Lambda-\mu)} \, . \tag{4}$$

The thermodynamic long wavelength behavior follows from the small momentum ($\mu \to 0$) behavior of the renormalized coupling constant. For $\left(g_{rs}^{\Lambda}\right)^2 < g_c^2 \equiv 3T^3/(4\pi^4\hbar^2 m v_F^2 \Lambda^4 S_4)$, the coupling constant is finite and the system is in the weakly interacting metallic phase. The divergence at $g_{rs} = g_c$ identifies one of the conditions for the Mott transition: the divergence of the renormalized coupling constant. The Mott critical point occurs for $U_c^2 = 4\hbar^2\Lambda^4 g_c^2 = 3T^3/(\pi^4 m v_F^2)$.

The divergence of the renormalized coupling constant is not enough to uniquely fix the Mott critical point. The missing ingredient is the behavior of quasiparticle motion and dynamics. The quasiparticle dynamics follows from its Green's function, which in the metallic phase is $G(\vec{k},\omega_n) = (-i\omega_n + v_F k)^{-1}$, with fermionic Matsubara frequency $\omega_n = (2n+1)\pi T/\hbar$. The fermionic Green's function is renormalized by fluctuations of density and spin density modes as illustrated in Figure 5, which provide a self energy correction to the fermion propagator appropriate to the distance scale $1/\mu$



FIG. 5: Feynman diagrams representing the self energy corrections to the quasiparticle propagator due to density and spin density fluctuations. Straight lines represent fermions, zig-zag lines are spin density fluctuations and wavy lines are density fluctuations.

$$\begin{aligned} \Sigma_\psi(\omega_n,\vec{k}) \simeq i\omega_n \, g_{rs}^2 \frac{2\hbar^2 m v_F^2}{\pi^6 T^3} \, \Lambda^3(\Lambda-\mu) \, \Bigg[\frac{\pi^4}{48} \\ + \frac{U^2 m^2 v_F^4}{64\pi^6 T^4} \, S_8\Bigg] \\ S_8 = \sum_n \frac{1}{(2n+1)^8} = \frac{17\pi^8}{(315)2^8} \end{aligned} \tag{5}$$

leading to a renormalized quasiparticle propagator $G(\vec{k},\omega_n) \simeq (-i\omega_n + v_F k - \Sigma_\psi)^{-1}$. This renormalized propagator has standard form with a renormalized Fermi velocity $v_F^R(\mu)$ and a renormalized quasiparticle weight $Z(\mu)$

$$\begin{aligned} G(\vec{k},\omega_n) \simeq \; & Z(\mu) \frac{1}{\left[-i\omega_n + v_F^R(\mu)\,k\right]} \\ Z(\mu) \simeq \; & 1 - g_{rs}^2 \frac{m\hbar^2 v_F^2}{\pi^4 T^3} \, \Lambda^3(\Lambda-\mu) \, \Bigg[\frac{\pi^2}{24} \\ & + \frac{17 U^2 m^2 v_F^4}{(315)2^{13} T^4}\Bigg] \\ v_F^R(\mu) \simeq \; & v_F \, \Bigg\{1 - g_{rs}^2 \frac{m\hbar^2 v_F^2}{\pi^4 T^3} \, \Lambda^3(\Lambda-\mu) \, \Bigg[\frac{\pi^2}{24} \\ & + \frac{17 U^2 m^2 v_F^4}{(315)2^{13} T^4}\Bigg]\Bigg\} \, . \end{aligned} \tag{6}$$

At long wavelengths, $\mu = 0$, we see that the renormalized Fermi velocity can be zero for a critical value of g_{rs}

$$(g_{rs})^2 = g_c^2 \equiv \frac{\pi^4 T^3}{m\hbar^2 v_F^2 \Lambda^4 \left[\frac{\pi^2}{24} + \frac{17 U^2 m^2 v_F^4}{(315)2^{13} T^4}\right]} \, . \tag{7}$$

This corresponds to a Fermi surface instability as the electrons immobilize and get localized. This vanishing of the Fermi velocity is equivalent to a divergence of the renormalized mass as seem from $v_F^R(\mu) = \hbar k_F / m^R(\mu)$. Furthermore the quasiparticle weight $Z(\mu)$ vanishes at $\mu = 0$ together with the vanishing of the Fermi velocity. The two conditions for the Mott critical point, the vanishing of the renormalized Fermi velocity and divergence of the renormalized coupling constant, need not be compatible. Making them consistent uniquely fixes the Mott critical point to occur at $T_c \propto U_c \propto m v_F^2$.

If we fix the temperature and coupling constant to their critical values, the renormalized Fermi velocity and spectral weight at momentum scale k are $Z(k) \simeq k/\Lambda$ and $v_F^R(k) \simeq v_F\, k/\Lambda$ leading to a quasiparticle propagation amplitude $G(\vec{k},\omega_n) \simeq k/\left[\Lambda(-i\omega_n + v_F k^2/\Lambda)\right]$ *Therefore we predict that at the novel Mott critical point recently measured[5], quasiparticle dispersion as measured via photoemission experiments will correspond to quadratically dispersing gapless fermions.* In fact, the retarded Green's function to be extracted from photoemission is

$$G(\vec{k},\omega) = i\frac{1}{\omega + i\delta - v_F\frac{k^2}{\Lambda}}\,\frac{1}{e^{\hbar\beta v_F k^2/\Lambda}+1} \,. \qquad (8)$$

The picture of the Mott transition that we have just obtained is that of metallic fermionic quasiparticles linearly dispersing perpendicular to the Fermi surface, which morph into gapless quadratically dispersing fermions which open a gap continuously as the system is tuned to the Mott insulating side. Furthermore $Z(0) \sim T - T_c$ and $Z(0) \sim U - U_c$ near the Mott critical point. *We thus also predict that the spectral weight at the Fermi level as measured by photoemission will go to zero linearly in the temperature deviation from criticality.*

One of the measured[5] novel exponents was that the deviation of the conductance from the critical conductance scales linearly with $T - T_c$. The conductivity of the system can be obtained from density or current fluctuations, which follow from the fermionic quasiparticles susceptibility via linear response theory

$$\sigma(\omega,\vec{k}) \sim \frac{\omega_n}{k^2}m\left[1 - \frac{|\omega_n|}{\sqrt{(v_F k)^2 + \omega_n^2}}\right] \,. \qquad (9)$$

The non-dynamical conductivity or conductance is given by the long wavelength and small frequency limits of the dynamical conductivity. The scattering rate, or inverse mean free time, introduces a frequency scale $\omega \sim 1/\tau$ below which the physics described by the conductivity is vitiated and smaller frequencies are unavailable. Similarly, the mean free path introduces a momentum scale $k \sim 1/l$. The conductance is then

$$G = \sigma(1/\tau, 1/l) \sim \frac{l^2}{\tau}m\left[1 - \frac{1}{\sqrt{(v_F\tau/l)^2 + 1}}\right] \,. \qquad (10)$$

In general the conductance is the sum of two conductances: that originating from scattering of impurities and that originating from scattering of phonons. Since impurities are fixed in space, such scattering has a fixed mean free path l, with scattering time subservient to l via $\tau = l/v_F$. The contribution to the conductance from impurities has the form $G_{imp} \sim v_F m$. Since the renormalizations of v_F and m cancel among themselves, this conductance is both constant in the metallic phase and at the Mott critical point. There is also a conductance arising from phononic scattering. Scattering from phonons, being a dynamical process, introduces an intrinsic scattering rate $1/\tau$ with the mean free path enslaved to it via $l = v_F\tau$. Therefore, the conductance arising from phonon scattering has the form $G_{phn} \sim v_F^2 m\tau$, which vanishes at the critical point as $T - T_c$.

Since the total conductance is $G = G_{imp} + G_{phn}$, the critical conductance is constant, arising from impurity scattering and thus nonuniversal. *We predict the critical conductance to be material dependent.* The deviations from the critical conductance for temperatures near T_c arise from phonon scattering. These deviations from the critical conductance vanish linearly with $T - T_c$ in disagreement with conventional LGW wisdom, but in agreement with experiment[5].

We thus conclude that our development can account correctly for at least one of the non-LGW exponents and thus points toward *the Mott critical point corresponding to a new universality class that, despite being finite temperature, is controlled by an essentially quantum aspect, fermionic fluctuations, and falls out of the reach of the Ginzburg-Landau-Wilson paradigm.*

[1] Chakravarty, S., Halperin B. I. & Nelson, D. R., Phys. Rev. B **39**, 2344 (1989).

[2] Damascelli, A., Hussain, Z. & Shen, Z.-X., Rev. Mod. Phys. **75**, 473 (2003).

[3] Stewart, G. R., Rev. Mod. Phys. **56**, 755 (1984).

[4] Joynt, R. & Taillefer L., Rev. Mod. Phys. **74**, 235 (2002).

[5] Kagawa, F., Miyagawa, K. & Kanoda K., Nature **436**, 534 (2005).

[6] Wilson, K. G., Rev. Mod. Phys. **47**, 773 (1975).

[7] Hertz, J. A., Phys. Rev. B **14**, 1165 (1976).

[8] Castellani, C., di Castro, C., Feinberg, D. & Ranninger, J., Phys. Rev. Lett. **43**, 1957 (1979).

[9] Kotliar, G., Lange, E. & Rozenberg. M. J., Phys. Rev. Lett. **84**, 5180 (2000).

[10] D. Bohm and D. Pines, Phys. Rev. **82**, 625 (1951); Phys. Rev. **85**, 338 (1952); Phys. Rev. **92**, 609 (1953); M. Gell-Mann and K. A. Brueckener, Phys. Rev. **106**, 364 (1957).